

Distinct Bacterial Consortia Established in ETBE-Degrading Enrichments from a Polluted Aquifer

Martina Kyselková, Joana Salles, Alain Dumestre, Yves Benoit, Geneviève Grundmann

► **To cite this version:**

Martina Kyselková, Joana Salles, Alain Dumestre, Yves Benoit, Geneviève Grundmann. Distinct Bacterial Consortia Established in ETBE-Degrading Enrichments from a Polluted Aquifer. Applied Sciences, MDPI, 2019, 9 (20), pp.4247. 10.3390/app9204247 . hal-02404233

HAL Id: hal-02404233

<https://hal-ifp.archives-ouvertes.fr/hal-02404233>

Submitted on 11 Dec 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Article

Distinct Bacterial Consortia Established in ETBE-Degrading Enrichments from a Polluted Aquifer

Martina Kyselková^{1,2,*}, Joana Falcão Salles^{1,3} , Alain Dumestre⁴, Yves Benoit⁵ and Geneviève L. Grundmann¹

¹ Université Lyon1, CNRS, UMR5557, Ecologie Microbienne, INRA, 69622 Villeurbanne CEDEX, France; j.falcao.salles@rug.nl (J.F.S.); Genevieve.Grundmann@univ-lyon1.fr (G.L.G.)

² Biology Centre of the Czech Academy of Sciences, Institute of Soil Biology, 37005 České Budějovice, Czech Republic

³ University of Groningen, Groningen Institute for Evolutionary Life Sciences, 9747 AG Groningen, The Netherlands

⁴ SERPOL, 69200 Vénissieux, France; alain.dumestre@serpol.fr

⁵ IFP Energies Nouvelles, 92852 Rueil-Malmaison, France; yves.benoit@ifpen.fr

* Correspondence: martinak@upb.cas.cz; Tel.: +33-387-775-770

Received: 16 August 2019; Accepted: 2 October 2019; Published: 11 October 2019



Abstract: Ethyl tert-butyl ether (ETBE) is a gasoline additive that became an important aquifer pollutant. The information about natural bacterial consortia with a capacity for complete ETBE degradation is limited. Here we assess the taxonomical composition of bacterial communities and diversity of the *ethB* gene (involved in ETBE biodegradation) in ETBE-enrichment cultures that were established from a gasoline-polluted aquifer, either from anoxic ETBE-polluted plume water (PW), or from an upstream non-polluted water (UW). We used a 16S rRNA microarray, and 16S rRNA and *ethB* gene sequencing. Despite the dissimilar initial chemical conditions and microbial composition, ETBE-degrading consortia were obtained from both PW and UW. The composition of ETBE-enrichment cultures was distinct from their initial water samples, reflecting the importance of the rare biosphere as a reservoir of potential ETBE degraders. No convergence was observed between the enrichment cultures originating from UW and PW, which were dominated by *Mesorhizobium* and *Hydrogenophaga*, respectively, indicating that distinct consortia with the same functional properties may be present at one site. Conserved *ethB* genes were evidenced in both PW and UW ETBE-enrichment cultures and in PW water. Our results suggest that the presence of *ethB* genes rather than the taxonomical composition of in situ bacterial communities indicate the potential for the ETBE degradation at a given site.

Keywords: ethyl tert-butyl ether; ETBE biodegradation; bacterial community; polluted aquifer; fuel oxygenates; *ethB* gene

1. Introduction

Oxygenates methyl tert-butyl ether (MTBE) and ethyl tert-butyl ether (ETBE) are widely used as enhancers of gasoline octane index, replacing previously used lead tetraethyl. These chemicals are, however, highly soluble in water and became a major pollutant in aquifers after accidental gasoline spills. Both compounds are poorly biodegradable and potentially carcinogenic. Their biodegradation often terminates with a toxic compound tert-butyl alcohol (TBA) [1,2]. The biodegradation of ETBE, in comparison to MTBE, has been studied less [3], but is of concern because ETBE has been found as an aquifer pollutant in some cases in Europe, especially in France [4].

Few bacterial isolates have been shown to degrade ETBE so far, belonging either to Betaproteobacteria, for example, *Aquicola tertiaricarbonis* L108, *Comamonas testosteroni* E1 and *Rubrivivax* sp. IFP 2047, or to Actinobacteria, for example, *Mycobacterium* sp. IFP 2009, *Rhodococcus equii* IFP 2002, *Rhodococcus zopfii* IFP 2005, *Rhodococcus ruber* IFP 2001, *Rhodococcus aetherivorans* IFP 2017, *Rhodococcus* sp. IFP 2042 and other strains, *Pseudonocardia tetrahydrofuranooxydans* K1 and *Pseudonocardia* sp. IFP 2050 [5–12]. The betaproteobacterial degraders were also able to degrade MTBE and TBA [6,8,12]. In all cases, the degradation of ETBE was observed only under aerobic conditions. In situ and microcosm studies, however, indicated that ETBE-degradation may also proceed under anaerobic conditions [13,14].

In *Rhodococcus ruber* IFP2001, it was shown that the operon *ethABCD* was essential for ETBE degradation [15]. The operon codes for a cytochrome P450 monooxygenase system involved in the initial oxydation of ETBE [1]. The expression of the gene *ethB*, encoding the cytochrome P450, is inducible by ETBE [16]. The *ethABCD* operon is flanked by class II transposon sequences, indicating that it may be horizontally transferred to other bacterial species. Indeed, highly conserved (>99% identity) *eth* genes were found in *Rhodococcus zopfii* IFP2005 and *Mycobacterium* sp. IFP2009 [7]. All these strains were able to grow on ETBE but accumulated TBA, showing that this operon itself is not sufficient to achieve complete ETBE mineralization.

The accumulation of TBA may present a problem for in situ treatments because of its toxicity. The co-culture with microorganisms able to grow on TBA may be required for successful bioremediation of oxygenate-contaminated sites. Indeed, a complete degradation of ETBE by a bacterial consortium was obtained from a polluted aquifer, in which *Rhodococcus* sp. degraded ETBE to TBA and *Bradyrhizobium* sp. further degraded TBA to CO₂ [13]. In addition, oxygenates in aquifers are usually accompanied by BTEX and other gasoline compounds, which implies that large bacterial consortia may be required for a full depollution of contaminated sites. Mentioned and other observations suggest the use of a community-based rather than a one-strain-based approach, as the best alternative to study these processes in the environment [17].

Few studies evaluated the composition of microbial consortia presenting oxygenate degradation. They indicated that the potential for intrinsic oxygenate degradation was dependent on the composition of the indigenous aquifer community, in combination with site-specific geochemical conditions [18–20]. These studies mainly focused on MTBE, while the information on the composition of microbial communities in ETBE polluted sites and natural consortia capable for ETBE biodegradation remain scarce [11,12,21,22]. In a study on bacterial enrichment cultures from five different aquifers, Le Digabel et al. [12] showed that the ETBE-degrading consortia had distinct taxonomical composition with prevailing Proteobacteria in three of them, and with Actinobacteria occurring in all of them. This confirmed the importance of the composition of the original aquifer communities, and also indicated which taxa might play a crucial role in ETBE degradation. Indeed, several actinobacterial and betaproteobacterial strains capable for partial or full ETBE degradation were isolated from these enrichment cultures [12]. The recent study of van der Walls et al. [22] has shown that algae (*Scenedesmus*, *Chlorella*) may be a part of natural microbial consortia from contaminated groundwater and, under light conditions, supply oxygen needed for bacterial ETBE degradation. They identified Comamonadaceae and Candidatus Moranbacteria as the most abundant bacterial groups in algal-bacterial culture that degraded ETBE under fed-batch conditions.

The present study was carried out in the context of the bioremediation of a gasoline-polluted aquifer in France, described by Fayolle-Guichard et al. [4]. Although BTEX have been already largely consumed in the plume at the time of sampling, ETBE remained as the prevailing pollutant, together with MTBE and TBA. Due to oxygen consumption in the course of BTEX biodegradation, the conditions in the plume became anoxic and therefore unfavorable for oxygenate biodegradation. Oxygenation of the aquifer is an option for ETBE bioremediation in such cases [4]. The main objective of this study was to establish aerobic ETBE-enrichment cultures from the studied aquifer that could later serve for bioaugmentation and assess their ETBE-degradation capacity and taxonomical composition. To this

end, the ETBE-enrichment cultures were established from water samples from the anoxic plume of the studied aquifer and from the upstream zone where the conditions were aerobic. Taxonomical composition of the original water samples and of the enrichment cultures was assessed with a 16S rRNA gene-based microarray, and the major taxa were confirmed by cloning and sequencing of the 16S rRNA genes. In addition, we assessed the presence and diversity of the *ethB* gene in original water samples and ETBE-enrichment cultures by cloning and sequencing.

2. Materials and Methods

2.1. Site Characteristics and Water Sample Collection

The experimental field was a leaking gas-station implemented with 40 piezometers, situated 10 km in the south of Lyon, Rhône region, France. The detailed geological, hydrodynamic and physico-chemical characterization of the field was done previously [4]. Water from two wells, one located at about 120 m upstream of the polluted zone and thus referred to as upstream well (UW; denoted Pz-1 in [4]) and another one within the polluted zone referred to as polluted well (PW; denoted P2 in [4]), was sampled for (i) bacterial community assessment (in June 2008, November 2008, March 2009 and April 2009) and (ii) the establishment of ETBE-enrichment cultures (in June 2008). The sampling depth was 2.7 and 1.9 meters for UW and PW, respectively. Water was first pumped out of the wells and one liter of water was then sampled after the wells were spontaneously re-filled. Water was kept at 4 °C until subsequent analyses (done on the next day). The chemical composition of water sampled in the two wells is indicated in Table 1.

Table 1. Chemical analysis of water from the upstream (UW) and the polluted (PW) well.

	ETBE (mg/L)	MTBE (mg/L)	TBA (mg/L)	BTEX (mg/L)	TOC ^a (mg/L)	Nitrates (mg/L)	Sulfates (mg/L)	Oxygen (mg/L)
PW	160.8	2.5	1.9	4.4	204	ND ^b	7	0.33
UW	ND	ND	ND	ND	ND	3	25	2.54

^a Total organic carbon, ^b ND—not detected.

2.2. Enrichment Cultures and Their Biodegradation Potential

Initial enrichment cultures and subsequent subcultures were carried out as follows (Figure S1). ETBE-enrichment samples are labeled I1 to I3 for the initial cultures and S1 to S3 for the subcultures. UW ETBE-enrichment subcultures S1–S3 were obtained from the initial UW culture I1, whereas PW ETBE-enrichment subcultures S1–S3 were obtained from the initial PW culture I3 (Figure S1). The establishment and handling of enrichment cultures was done in a laminar flow hood to maintain sterile conditions. For initial enrichment cultures, water samples (20 mL) from UW and PW were filtered (0.22 µm) to discard any pollutant that could be dissolved in water. The filters were introduced to air-tight 500 mL-Schott flasks containing 150 mL mineral medium (KH₂PO₄ 1.4 g/L; K₂HPO₄ 1.7 g/L; MgSO₄ × 7 H₂O 0.5 g/L; NH₄NO₃ 1.5 g/L; CaCl₂ × 2 H₂O 0.04 g/L; FeSO₄ × 7 H₂O 1 mg/L) with 100 mg/L ETBE representing the sole source of carbon. A vitamin solution and an oligo-element solution were added as previously described [23]. The headspace in the flasks was calculated so that oxygen was in a large excess with regard to the theoretical oxygen demand of ETBE. The initial enrichment cultures were set in triplicates and incubated under agitation at 30 °C. The flasks were closed with a stopper and equipped with a side-arm enabling sampling in the course of incubation. Samples of 2 mL were regularly withdrawn to measure ETBE and TBA concentrations (see further). When ETBE was fully consumed, the cultures were spiked with ETBE so as to reach the initial concentration (1 to 3 times). The initial enrichment cultures were sampled (40 mL) at 67 days (replicate I3 of PW initial enrichment culture) and 253 days (the other replicates), centrifuged, and cell pellets were stored at −80 °C for DNA extraction.

After the complete biodegradation of ETBE in the initial enrichment cultures, triplicate enrichment subcultures were established from one flask for UW (I1) and PW (I3), by transferring 30 mL of the initial enrichment cultures into 150 mL fresh minimal medium containing 100 mg/L ETBE (Figure S1). After 1 week, 40 mL of the enrichment subcultures were centrifuged and the bacterial pellets were stored at $-80\text{ }^{\circ}\text{C}$ until DNA extraction. After the complete ETBE biodegradation, the enrichment subcultures were spiked with ETBE so as to reach the initial concentration and the oxygenate concentrations were further monitored. This was done twice in order to confirm the biodegradation capacities of the established community.

ETBE and TBA were quantified by flame ionization detection on a Varian 3300 gas chromatograph (Varian, Les Ulis, France) equipped with a $0.32\text{ mm} \times 25\text{ m}$ Porabond-Q capillary column (J&W Scientific, Chromoptic, Auxerre, France), using a two-step temperature gradient ranging from $105\text{ }^{\circ}\text{C}$ to $210\text{ }^{\circ}\text{C}$ at $10\text{ }^{\circ}\text{C}/\text{min}$. The column was then maintained at $210\text{ }^{\circ}\text{C}$ for 20 min. Helium ($1.6\text{ mL}/\text{min}$) was used as the carrier gas. Samples were filtered through $0.22\text{ }\mu\text{m}$ filters (Prolabo, Fontenay-sous-Bois, France) and injected without further treatment.

2.3. DNA Extraction

One liter of water collected from the wells was filtered through sterile $0.2\text{ }\mu\text{m}$ nylon filter (Millipore, Molsheim, France) and water DNA was extracted from the filters with UltraClean Water DNA kit (MoBio Laboratories, Carlsbad, CA, USA) according to the manufacturer's protocol.

The enrichment cultures (notably those originating from UW) contained visible flocks suggesting the presence of actinomycetes. Therefore, a modified method of Pospiech and Neumann [24] was used for total DNA extraction from all enrichment cultures. Bacterial pellets from 40 mL of enrichment cultures were resuspended in $162\text{ }\mu\text{L}$ SET buffer (75 mM NaCl , 25 mM EDTA , 20 mM Tris , pH 8.0) with lysozyme (Sigma, L'Isle d'Abeau, France; $1\text{ mg}/\text{mL}$) and transferred to 1.5 mL Eppendorf tubes. Immediately, $16\text{ }\mu\text{L}$ of lysostaphin (Sigma; $50\text{ }\mu\text{g}/\text{mL}$ in 10 mM Tris-HCl , pH 7.5) were added to each tube and the tubes were incubated at $37\text{ }^{\circ}\text{C}$ for 1 h. The remaining flocks were disintegrated by forcing the samples through 0.4 mm diameter syringe needles. After addition of $20\text{ }\mu\text{L}$ SDS (10%) and $12\text{ }\mu\text{L}$ proteinase K ($23\text{ mg}/\text{mL}$; NucleoSpin Tissue kit, Macherey-Nagel, Hoerd, France), the samples were incubated at $55\text{ }^{\circ}\text{C}$ for 2 h. DNA extraction and purification was carried out with NucleoSpin Tissue kit (Macherey-Nagel, Hoerd, France) from step 3 according to the manufacturer's instructions. DNA concentrations were assessed using Nanodrop ND-100 spectrophotometer (Labtech, Palaiseau, France).

2.4. PCR Amplification of 16S rRNA (*rrs*) Genes and *ethB* Genes

The universal eubacterial primers T7-pA (forward; *TAATACGACTCACTATAGAGAGTTTGATCCTGGCTCAG*) and pH (reverse; *AAGGAGGTGATCCAGCCGCA*) were used to amplify almost full-length 16S rRNA gene [25]. Primer T7-pA includes the sequence of T7 promoter (in italics) at the 5' end, which enabled T7 RNA polymerase-mediated in vitro transcription using the PCR products as templates. PCR reactions were carried out in a total volume of $50\text{ }\mu\text{L}$ and the reaction mixture prepared as in [26].

The *ethB* genes from UW and PW water samples (sampled in June 2008), and from UW ETBE-enrichment subculture S1 and PW ETBE-enrichment subculture S3 (Figure S1) were amplified with primers *ethB*-F2 (forward; *CACGCGCTCGGCGACTGGCAGACGTTTCAGT*) and *ethB*-R2 (reverse; *TCCGACGCACATGTGCGGGCCGTACCCGAA*) [19]. The primers were defined based on known *ethB* sequences from *Mycobacterium* sp. IFP2009, *Rhodococcus zopfii* IFP2005 and *Rhodococcus ruber* IFP2001 [7]. The primers anneal to the *Mycobacterium* sp. IFP2009 *ethB* sequence positions 175 and 1027, yielding a 897 bp-long fragment. PCR reactions were carried out in $50\text{ }\mu\text{L}$ total volume. PCR reaction mixtures contained $1 \times$ reaction buffer for Taq DNA Polymerase (Invitrogen, Cergy Pontoise, France), 1.5 mM MgCl_2 (Invitrogen, Cergy Pontoise, France), $0.5\text{ }\mu\text{M}$ of each primer (Invitrogen, Cergy Pontoise, France), $50\text{ }\mu\text{M}$ of each dNTP, 20 ng of template DNA, 0.025 mg of T4 gene 32 protein (Roche, Meylan, France), 2% dimethyl sulfoxide and 10 U of Taq DNA Polymerase (Invitrogen, Cergy Pontoise, France). Negative controls (sterile ultra-pure water pipetted to PCR reactions instead of template DNA) and

positive controls (20 ng of *Rhodococcus ruber* IFP 2001 genomic DNA as template) were always included. Thermal cycling was carried out with an initial denaturation step of 95 °C for 7 min, followed by 30 cycles with 30 s denaturation at 94 °C, 30 s annealing at 68 °C, 60 s elongation at 72 °C, and a final elongation step for 7 min at 72 °C. When no visible PCR products were obtained (water samples), the PCR was repeated using 7 µL from the previous PCR mix as a template. PCR products were purified with MinElute PCR purification kit (Qiagen, Hilden, Germany) and DNA concentrations were assessed with Nanodrop ND-100 spectrophotometer (Labtech, Palaiseau, France).

2.5. Sequencing and Sequence Analysis

Purified *rrs* PCR products obtained from UW and PW water sampled in June 2008, and from UW ETBE-enrichment subculture S1 and PW ETBE-enrichment subculture S3 (Figure S1), and *ethB* PCR products obtained from PW water (June 2008) and UW ETBE-enrichment subculture S1 and PW ETBE-enrichment subculture S3 were cloned into the cloning vector pGEM-T (pGEM-T Easy Vector System kit; Promega, Charbonnières, France) according to the manufacturer's protocol. For *rrs* genes, 96 clones per sample were sequenced on both strands using Sanger sequencing (AGOWA, Berlin, Germany). For *ethB* genes, 48 clones per sample were sequenced on one strand (AGOWA). Sequences were checked and edited with BioEdit version 5.0.9 (<http://www.mbio.ncsu.edu/BioEdit/BioEdit.html>). Chimeric 16S rRNA gene sequences were identified using the chimera detection program Bellerophon [27]. The good quality sequences of *rrs* (63, 65, 66 and 74 sequences from PW water, UW water, PW ETBE-enrichment subculture S3 and UW ETBE-enrichment subculture S1, respectively) and *ethB* (23, 42 and 45 sequences from PW ETBE-enrichment subculture S3, UW ETBE-enrichment subculture S1 and PW water, respectively) are available at GenBank (accession numbers KP636042-KP636419).

Sequence affiliation was performed using RDP Classifier [28] at Ribosomal RDP Release 11 (<http://rdp.cme.msu.edu/index.jsp>; Update 5, 30 September 2016) and algorithm BlastN with default parameters at NCBI Blast (<http://www.ncbi.nlm.nih.gov/blast/Blast.cgi>). Phylogeny reconstructions were done with MEGA4 software version 4.0.2 (Center for Evolutionary Medicine and Informatics, Tempe, AZ) [29], based on sequence alignments done with ClustalW version 1.83 (Conway Institute UCD, Dublin, Ireland), using default settings. Neighbor-joining trees were built based on phylogenetic distances calculated with Kimura 2 parameter [30,31]. Diversity indices (Shannon index and Chao1 richness estimator) were calculated with RAMI software (Björn Canbäck Bioinformatics, Lund, Sweden) [32].

2.6. Definition of New Probes for 16S rRNA Gene-Based Taxonomical Microarray

Sixty-eight new 16S rRNA probes were defined (Table S1) and added to the previous probe set [33] containing 1033 probes targeting 19 bacterial phyla at different taxonomic levels in order to improve microarray detection potential of bacteria associated to oxygenate pollution. The probes (20mers) were designed using ARB software (<http://www.arb-home.de>) [34], based on 16S rRNA sequences recovered from the aquifer and the enrichment cultures (see above) as well as 16S rRNA sequences of known ETBE and MTBE degrading species (e.g., *Aquicola tertiarycarbonis*, *Variovorax paradoxus*, *Rhodococcus aetherivorans*, *Rhodococcus ruber*, *Mycobacterium austroafricanum* and *Pseudonocardia* spp.), using the previously defined parameters of the Probe Design function [26,35]. The probe specificity was tested with the Probe Match function against the reference Silva-94 database [36], and only the probes that displayed weighted mismatch value above 1.5 with the non-targeted taxa [33] were retained. The hybridization properties of probes (e.g., melting temperature, potential formation of secondary structures and 3' dimers) were further tested in silico, according to [26,35].

2.7. In Vitro Transcription, Microarray Hybridization, Scanning and Image Analysis

In vitro transcription of 16S rRNA genes was performed on the purified PCR products (400 ng) to obtain fluorescently (Cy3) labeled 16S rRNA for microarray hybridization [37]. This procedure

and 16S rRNA fragmentation are described in [35]. Labeled samples were hybridized to the 16S rRNA gene-based taxonomic microarray (containing 1033 probes from [33] and 68 new probes). This microarray has been previously validated by 16S rRNA gene sequencing and qPCR [33,38]. Each microarray probe was present in four copies per slide. Hybridization and image analysis were carried out according to [35].

2.8. Microarray Data Filtration and Statistical Analysis

Data filtration and normalization was conducted as in [33]. Differences in bacterial community composition were statistically assessed with analysis of similarities (ANOSIM) using Primer6 software (PRIMER-E Ltd., Plymouth, UK) [39]. Distances between the groups of samples were analyzed by multidimensional scaling (MDS; Primer6, based on Bray-Curtis similarity matrix) [40]. The probe contribution to the differences between significantly different groups was assessed by similarity percentage analysis (SIMPER; Primer6). Probes giving the cumulative contribution up to 50% were retained and their significance for distinguishing between the groups was further tested with Wilcoxon rank sum tests ($P < 0.05$), using R (<http://www.r-project.org>).

3. Results

3.1. Bacterial Communities in Polluted and Upstream Well Water

MDS (Figure 1) and ANOSIM on microarray data showed that bacterial communities in UW and PW water significantly differed ($P < 0.05$) (the four sampling dates were grouped together). There were 99 positive probes in common between UW and PW water, 99 probes were positive only in UW water and 41 were positive only in PW water. SIMPER analysis together with Wilcoxon rank sum tests showed which probes significantly ($P < 0.05$) contributed to up to 50% of the differences in bacterial community composition between UW and PW (Figure S2). Notably, the signal intensities from probes targeting sequences affiliated to Spirochates (trepncln), alphaproteobacterial genus *Azoarcus* (Arcus1, Arcus2, AZA463), betaproteobacterial genera *Nitrosospira* (Nit1A, Nit1C), *Acidovorax* (AcidP2cln, Acave3, Acave4), *Polaromonas* (Polarcln), and family Comamonadaceae (Rhodofcln; with highest identities to *Albidiferax*), and deltaproteobacterial genus *Desulfovibrio* (DVLVT222f) were higher in PW water than UW water. In contrast, signals from probes affiliated to the family Rhizobiaceae (Rzbc1247, Alphaproteobacteria), family Oxalobacteraceae (Dugacln, Betaproteobacteria), betaproteobacterial genera *Ralstonia/Cupriavidus/Wautersia* (Ralstcln2) and *Janthinobacterium* (Janaga3), *Pseudomonas* (Pseud; Gammaproteobacteria) and *Nitrosospira* (NSR1156 and NSR826) were higher in UW water than in PW water.

Bacterial communities from PW and UW water sampled in June 2008 were analyzed by 16S rRNA gene sequencing to confirm the presence of the distinguishing bacterial taxa between the two wells. The bacterial community in UW water was more diverse (Shannon index 2.97, 48 OTUs estimated with Chao1 at 97% level of sequence identity) than in PW water (Shannon index 2.61, 34 OTUs estimated) (Figure S3). The only known genus shared by UW and PW water, based on 16S rRNA gene sequencing, was *Opitutus* (Verrucomicrobia) (Figure S4). The UW water was dominated by Alphaproteobacteria (mainly *Sphingomonas* and *Methylocystis*), while sequences related to known anaerobes (e.g., epsilonproteobacterium *Arcobacter*) and Betaproteobacteria (unclassified Betaproteobacteria, Rhodocyclaceae and Comamonadaceae) prevailed in PW. This is concordant with microarray results. Microarray probes targeting these groups, that is, Sphingo4 and Sphingo5A for Sphingomonadaceae, Methylocyst1 for Methylocystaceae, Rhodofcln for sequences related to Comamonadaceae and Arco1 for *Arcobacter* were always positive when these taxa were found with the 16S rRNA sequencing approach, although they were not always revealed by SIMPER as the most contributing to the differences between UW and PW.




Figure 1. Multidimensional Scaling of bacterial communities from PW water (black circles), UW water (white circles), PW ETBE-enrichment cultures (black triangles) and UW ETBE-enrichment cultures (white triangles). Water samples are labeled with month/year of sampling. ETBE-enrichment samples are labeled I1–I3 for the initial cultures and S1–S3 the subcultures (see Figure S1 for more details). Samples that were used for 16S rRNA and *ethB* gene sequencing are marked with an asterisk. The distances between samples were based on Bray-Curtis similarity matrix, which was calculated from normalized probe signals. Low 2D stress value (<0.1) indicates that the distances between samples in reduced 2-dimensional space correspond well with the actual multivariate distance between the samples.

3.2. ETBE Biodegradation Potential in the ETBE Enrichments from the Upstream and the Polluted Well

Both UW and PW initial enrichment cultures degraded ETBE, though the initial enrichment cultures from UW showed heterogeneous biodegradation characteristics (i.e., the time after which the biodegradation occurred and whether they accumulated TBA in the course of ETBE degradation; Figure S1). One of the initial enrichment cultures from each well that showed a complete biodegradation of at least two doses of 100 mg/L ETBE, was then used to establish the enrichment subcultures to confirm the biodegradation capacities and to measure the ETBE biodegradation rates (Figure S1). The mean biodegradation rates in the UW and PW subcultures were 19.6 and 24.7 mg ETBE L⁻¹ × d⁻¹, respectively (Figure S5). TBA accumulated only transiently during ETBE biodegradation in UW subcultures (Figure S5A) and reached a maximum concentration of 53.9 mg/L during the first ETBE dose degradation. In contrast, TBA concentrations increased with time in all PW subcultures, reaching a maximum of 180.3 mg/L after the consumption of the third ETBE dose (Figure S5B).

3.3. Bacterial Communities in the ETBE-Enrichment Cultures

The microarray analysis of the ETBE-enrichment communities originating from UW and PW wells revealed that they differed significantly from the corresponding water communities as well as from each other (Figure 1; ANOSIM, $P < 0.01$ in all cases). There was no convergence between the ETBE-enrichment communities originating from UW and PW (Figure 1). The initial enrichment cultures and the subcultures originating from the same well grouped together, showing little turnover in the course of the experiment. Also, the three initial enrichment cultures from UW grouped together

(Figure 1), despite their distinct biodegradation rates and differences in their TBA biodegradation potential (Figure S1).

SIMPER analysis together with Wilcoxon rank sum tests showed which probes significantly ($P < 0.05$) contributed to up to 50% differences in bacterial community composition between UW- and PW-enrichment cultures (Figure S2). Notably, the enrichments (initial cultures and subcultures grouped together) from PW, in comparison to the UW enrichments, had higher signals for probes targeting Betaproteobacteria belonging to *Hydrogenophaga* (HyMa2, HydETBE1, HydETBE3), *Nitrosospira* (Nit1B, Nit1C), *Variovorax* (Varpar) and *Aquabacterium* (Aquacln2), Gammaproteobacteria *Nitrosococcus* (Nibmob) and *Eikenella/Neisseria* (Eikcor2), and *Paenibacillus* (Palv, Palv2; Firmicutes). In contrast, the UW enrichment cultures had higher signals for probes targeting Alphaproteobacteria from the family Rhizobiaceae (Rhizo157, Rhi) related to *Mesorhizobium/Rhizobium/Devosia* (Phyllobact, Mesocln1, Davocln1), from Actinobacteria *Pseudonocardia* (Psdonoca), *Gordonia* (Gordcln3) and *Mycobacterium* (Mycoba2), and from Chloroflexi (Dehacln5, Dehacln6), as compared to PW enrichments. These probes distinguishing the UW and PW enrichments also significantly contributed to the differences between the enrichments and their respective initial water samples (Figure S2). Though there were no overall differences in the community composition between the three initial ETBE enrichment cultures from UW (Figure 1), it is interesting to note that the enrichment culture I3, which showed only a slight ETBE biodegradation activity at the time of sampling, lacked signal from the probe Mesocln1 for *Mesorhizobium* (Figure S2).

There was no significant difference in bacterial community composition between the enrichments that completely degraded ETBE and those that degraded ETBE but accumulated TBA, based on ANOSIM analysis of microarray data. When individual probes were analyzed with Wilcoxon rank sum tests ($P < 0.05$), only XAX818 (for *Xanthomonas*) gave significantly higher signals in the enrichments that showed a complete ETBE degradation, while Ralstcln2 (for *Ralstonia/Cupriavidus/Wautersia*, including certain clones from UW) gave significantly lower signals.

The dominating taxa in the two ETBE-enrichment subculture communities originating from UW and PW that were able to degrade ETBE were further studied by sequencing the 16S rRNA gene. The diversity in the UW ETBE-enrichment subculture S1 was higher (Shannon index 2.23 and 26 estimated OTUs) than the diversity in the PW ETBE-enrichment subculture S3 (Shannon index 1.03, 7 estimated OTUs) (Figure S3). As far as could be concluded from the number of analyzed clones, no genus was shared by the two enrichment subcultures (Figure 2). According to the 16S rRNA gene-sequencing, the UW ETBE-subculture community was enriched in alphaproteobacterial taxa different from those found in the well water community, with prevailing *Mesorhizobium* clones (31% of all clones) (Figure 2). Phylogenetic analysis (Figure S6) showed that the putative *Mesorhizobium* strains were diverse, with one sequence close to *M. plurifarum*, two sequences close to *M. loti* et *M. ciceri* and 33 sequences clustering apart from the established *Mesorhizobium* species. Besides, the presence of Betaproteobacteria, Gammaproteobacteria, Actinomycetes, Chloroflexi and Nitrospira was confirmed by 16S rRNA gene-sequencing in the UW ETBE-enrichment subculture S1. The ETBE-enrichment subculture S3 from PW was dominated by Betaproteobacteria differing from those found in PW water, mainly by *Hydrogenophaga* (67% of all clones) (Figure 2). The *Hydrogenophaga* sequences, clustering apart from the established *Hydrogenophaga* species, showed little diversity (Figure S7) and the majority of sequences were identical to the *Hydrogenophaga* sp. Rs71 found in a BTEX-enrichment culture from a gasoline-contaminated site in the UK [41]. *Acidovorax* and three unknown clades of Betaproteobacteria (with the closest relatives identified by BLAST as *Hydrogenophaga*, *Cupriavidus* and *Ralstonia* at about 95% sequence identity levels), as well as a few Gammaproteobacteria and Acidobacteria were also found in the PW ETBE-enrichment subculture S3.

The prevailing taxa in the PW and UW ETBE-enrichment subcultures, according to 16S rRNA gene sequencing, were targeted by the probes HydETBE3 and HydETBE1 (for *Hydrogenophaga*) and Mesocln1 (for *Mesorhizobium*). Based on signals from these probes, *Hydrogenophaga* was also detected in UW and PW water, while *Mesorhizobium* was not detected in any water sample.




Figure 2. Comparison of PW and UW ETBE-enrichment subculture communities by 16S rRNA gene sequencing (based on 66 and 74 clones, respectively). Bars represents per cents of recovered clones from PW (black) and UW (white) ETBE-enrichment subcultures affiliated to the indicated bacterial taxa.

3.4. Genetic Diversity of the *ethB* Gene

The *ethB* gene primers we used generated a positive signal in PW water and in ETBE-enrichment subcultures originating from both UW and PW. In PW water, however, the PCR product was obtained only after re-amplification of the first PCR reaction, indicating that the abundance of *ethB* targeted by these primers in PW water was low. All efforts to amplify *ethB* from UW water samples were unsuccessful. All obtained *ethB* sequences shared about 99% identity. Based on nucleotide blast, they were highly similar (99% identity) to *ethB* genes of *R. ruber* IFP2007 (accession number FJ481920), *R. zopfii* (FJ481919), *Rhodococcus* sp. PEG604 (DQ847178), *Mycobacterium* sp. IFP2009 (FJ481921) and

Aquicola tertiarycarbonis L108 (KC333876), and had a high G+C content (about 68%). The *ethB* sequences recovered from UW ETBE-enrichment subculture S1 could be distinguished by phylogenetic analysis from those recovered from PW ETBE-enrichment subculture S3, but the bootstrap value was low (i.e., 47; Figure 3). The sequences obtained from PW water clustered together either with UW ETBE-enrichment subculture S1 sequences or with PW ETBE-enrichment subculture S3 sequences or separately. However, none of the sequences recovered from the ETBE-enrichment subcultures was 100% identical to those recovered from PW water.




Figure 3. Phylogenetic tree based on *ethB* sequences recovered from PW ETBE-enrichment subculture S3 (23 sequences), UW ETBE-enrichment subculture S1 (42 sequences) and PW water (45 sequences), and highly similar (>99% identity, according to blastn) *ethB* sequences from published strains. The Neighbor-Joining tree was constructed based on phylogenetic distances calculated with Kimura-2 parameter (816 nucleotide positions included in the analysis), and bootstrapped over 1000 replicates. The collapsed parts of the tree (presented as black triangles) are labeled with the corresponding samples and numbers of clones (in brackets). Bootstrap values are marked in bold next to the branches. The scale bar shows phylogenetic distance.

4. Discussion

We used aerobic enrichment cultures with ETBE as sole source of carbon to reveal the potential for aerobic ETBE degradation at the studied aquifer, and to identify ETBE-degrading consortia. Water from two wells, one from the anoxic plume of the aquifer, and one situated upstream the polluted zone were used to establish the enrichment cultures. Distinct consortia with ETBE-biodegradation potential could establish from the two wells. Moreover, the *ethB* gene was distributed independently of the communities' taxonomic composition. From an ecological perspective, these results highlight the importance of the so-called rare biosphere as a potential reservoir of ETBE degraders.

This work has been done in a context of a large bioremediation study on a gasoline-polluted aquifer, where bacterial communities and chosen biodegradation genes were monitored in situ over time, using a high-throughput taxonomic microarray and PCR of functional genes (not shown). The microarray has been previously developed to target 19 bacterial phyla at different taxonomic levels [34,39], and it was implemented with 68 new probes targeting uncultured bacteria from the studied aquifer as well as known ETBE and MTBE degrading isolates. The microarray has been previously validated with sequencing and qPCR. It has been shown that the microarray was semi-quantitative, enabling between-sample comparisons based on signal intensities from individual probes [33,38,42]. Between-probe comparisons (i.e., between-taxon comparisons) are, however, not quantitative. Therefore, we used cloning-sequencing of chosen samples to reveal the main taxa present. Though both methods bring some bias due to PCR, there was a good agreement between the two methods concerning the presence of bacterial taxa. Not surprisingly, microarray revealed some taxa that were not shown by cloning-sequencing, which is consistent with its higher sensitivity [42].

This study has shown that the bacterial community in the polluted well, compared to the upstream well, had a reduced diversity, which is consistent with previous studies on BTEX contaminated sites [41,43]. It was enriched in Betaproteobacteria and in anaerobe-like taxa, in line with the anaerobic character of the well (Table 1). Similar results were obtained with water from other wells situated in the plume (not shown). The depletion of oxygen in BTEX-contaminated sites is a common state and the redox potential seems to be the main factor driving the composition of bacterial communities in such sites [44]. Despite the distinct initial conditions, ETBE-degrading consortia could be established from water of both aerobic unpolluted well and the anaerobic polluted well.

Comparison of the original aquifer communities with their respective enrichment cultures showed that the biodegradation consortia might represent only a minor part of the original aquifer community. Deeper techniques such as next-generation sequencing could be helpful to estimate the relative abundance of the rare taxa in original aquifer samples. The failure of biostimulation approach (either oxygen or oxygen and salt addition) in further laboratory experiments with polluted water from the studied aquifer [4] confirms the scarcity of the ETBE-degrading bacteria. In contrast, bioaugmentation using the UW ETBE-enrichment culture characterized in this study led to the complete ETBE biodegradation of polluted water under laboratory conditions at 15 °C, which is the mean aquifer temperature [4].

The taxonomical composition of the enrichment communities did not reflect their biodegradation potential. On the one hand, the initial ETBE-enrichment cultures from the UW had very similar taxonomical composition, while they differed in their capacity of ETBE and TBA degradation. On the other hand, enrichment cultures from UW and PW that degraded ETBE were composed by completely distinct bacterial taxa. These results highlight the high degree of functional redundancy of the aquifer and enrichment communities. The lack of convergence in bacterial community composition in contaminated aquifers or soils was also noted in other studies [44,45], indicating that distinct consortia with the same functional properties may be present at one site. Consequently, similarities in taxonomical composition of ETBE-degrading bacterial consortia from different studies and sites can be seen only on high taxonomic levels. Alpha-, Beta- and Gammaproteobacteria, Actinobacteria, Acidobacteria and Chloroflexi are reported in this and other studies [12,22], prompting to search candidate biodegrading bacteria within these groups.

We did not aim to test individual strains from ETBE-enrichment cultures for their biodegradation potential, it is worth noting, however, that several bacterial genera that include strains with ETBE-biodegradation properties such as *Mycobacterium*, *Pseudonocardia* and *Variovorax* were detected in the ETBE-enrichment cultures by microarray. *Pseudonocardia* was additionally confirmed in UW ETBE-enrichment culture by sequencing. It is not known whether the *Mesorhizobium* and *Hydrogenophaga* spp. dominating in the UW and PW ETBE-enrichment clone libraries, respectively, were involved in the initial ETBE oxygenation step or in the degradation of other compounds occurring in the course of ETBE biodegradation, such as TBA. A slight ETBE- (and no TBA-) degradation capacity was reported once in an isolate of *Mesorhizobium* from a bacterial enrichment culture degrading mixture of hydrocarbons and gasoline additives [46]. *Mesorhizobium* spp. were also detected in three out of five ETBE-enrichments originating from different aquifers in the study of Le Digabel et al. [12], and once they even reached the relative abundance of 22%, but no ETBE-degrading strain was obtained in their study. ETBEdegradation was never reported in *Hydrogenophaga*, although one *Hydrogenophaga* strain (ENV735) was shown to be able to degrade MTBE and TBA [47]. *Hydrogenophaga* was suggested to have the main role in MTBE biodegradation in the presence of BTEX [48], while it was considered only as an accompanying microflora in a MTBE-enrichment culture from a gasoline-polluted site [49]. In our study, *Hydrogenophaga* was enriched from the BTEX-polluted plume of the aquifer and its ability to degrade oxygenates merits further studies.

The successful amplification of *ethB* genes from both types of enrichments as well as from the polluted well confirms that the gene *ethB* is a good candidate for an in situ marker of the ETBE biodegradation potential, as suggested in [4]. The great level of conservation observed for the *ethB*

gene obtained from highly distinct bacterial communities and consortia noticed in this study further indicates that this gene may be horizontally transferred among different hosts. The natural diversity of *ethB* may, however, remain hidden as the currently used *ethB* primers [7] are based on only three sequences. Indeed, van der Waals et al. [22] did not detect *ethB*, when using primers based on *A. tertiarycarbonis* L108 sequence [50], in their ETBE-degrading cultures from polluted groundwater, despite the fact that ETBE-degradation was shown to be cytochrome P450-dependent. Besides the possibility that other than EthABCD system was involved, it is also conceivable that the primers used had not enough sequence complementarity with the *ethB* gene present. We suggest that *ethB* gene diversity, abundance and expression should be taken into consideration when evaluating in situ potential for ETBE degradation.

5. Conclusions

The functional redundancy of ETBE-degrading bacterial consortia observed in this work shows that the taxonomical composition provides little information regarding potential for the ETBE degradation at a given site, while the gene *ethB* appears to be a good indicator of ETBE-degradation potential for further in situ operations.

Supplementary Materials: The following are available online at <http://www.mdpi.com/2076-3417/9/20/4247/s1>, Figure S1: A simplified overview of the ETBE-enrichment cultures from PW and UW water and their biodegradation properties, Figure S2: Microarray comparisons of bacterial communities between PW and UP water, PW water and PW ETBE-enrichments, UP water and UP ETBE-enrichments, and PW ETBE-enrichments and UW ETBE-enrichments, Figure S3: Analysis of the 16S rRNA gene diversity in UW and PW water and ETBE-enrichment subcultures based on cloning-sequencing, Figure S4: Comparison of UW and PW water bacterial communities by 16S rRNA gene sequencing, Figure S5: Degradation of ETBE and TBA in UW ETBE-enrichment subcultures S1–S3 and PW ETBE-enrichment subcultures S1–S3, Figure S6: Phylogeny analysis of 16S rRNA clones from the UW ETBE-enrichment subculture S1, Figure S7: Phylogeny analysis of 16S rRNA clones from the PW ETBE-enrichment subculture S3, Table S1: List of new microarray probes designed in this study and their targets.

Author Contributions: Conceptualization, M.K., J.F.S., A.D., Y.B. and G.L.G. Methodology, M.K., J.F.S., Y.B. and G.L.G.; Validation, M.K., Y.B. and G.L.G.; Formal Analysis, M.K.; Investigation, M.K., J.F.S. and Y.B.; Resources, A.D., Y.B. and G.L.G.; Data Curation, M.K.; Writing—Original Draft Preparation, M.K. and G.L.G.; Visualization, M.K.; Supervision, A.D. and G.L.G.; Project Administration, A.D.; Funding Acquisition, A.D.

Funding: This work was funded by the project TISATIE (*Traitements in situ alternatifs: Traçabilité, Innocuité, Efficacité: Application aux polluants pétroliers type MTBE, ETBE*), funded by VALORSITES AXELERA, a French competitiveness cluster.

Acknowledgments: We thank Jacqueline Haurat (CNRS, UMR5557, Écologie Microbienne; Villeurbanne; France), and Jean Philippe Tracol (SERPOL; Vénissieux, France) and SERPOL technical team for technical help.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Lopes Ferreira, N.; Malandain, C.; Fayolle-Guichard, F. Enzymes and genes involved in the aerobic biodegradation of methyl *tert*-butyl ether (MTBE). *Appl. Microbiol. Biot.* **2006**, *72*, 252–262. [[CrossRef](#)] [[PubMed](#)]
2. Rosell i Linares, M. Vigilancia Ambiental de Metil *Tert*-Butil eer (MTBE) un Additiu de la Gasolina en Aigües i Sols. Ph.D. Thesis, Universitat de Barcelone Facultat de Chimie, Barcelona, Spain, July 2006.
3. Hyman, M. Biodegradation of gasoline ether oxygenates. *Curr. Opin. Biotechnol.* **2013**, *24*, 443–450. [[CrossRef](#)] [[PubMed](#)]
4. Fayolle-Guichard, F.; Durand, J.; Cheucle, M.; Rosell, M.; Michelland, R.J.; Tracol, J.P.; Le Roux, F.; Grundmann, G.; Atteia, O.; Richnow, H.H.; et al. Study of an aquifer contaminated by ethyl *tert*-butyl ether (ETBE): Site characterization and on-site bioremediation. *J. Hazard. Mater.* **2012**, *201*, 236–243. [[CrossRef](#)] [[PubMed](#)]
5. Fayolle, F.; Hernandez, G.; Le Roux, F.; Vandecasteele, J.P. Isolation of two aerobic strains that degrade efficiently ethyl *t*-butyl ether (ETBE). *Biotechnol. Lett.* **1998**, *20*, 283–286. [[CrossRef](#)]

6. Kharoune, M.; Kharoune, L.; Lebeault, J.M.; Pauss, A. Isolation and characterization of two aerobic bacterial strains that completely degrade ethyl tert-butyl ether (ETBE). *Appl. Microbiol. Biot.* **2001**, *55*, 348–353. [[CrossRef](#)] [[PubMed](#)]
7. Béguin, P.; Chauvaux, S.; Miras, I.; Francois, A.; Fayolle, F.; Monot, F. Genes involved in the degradation of ether fuels by bacteria of the *Mycobacterium/Rhodococcus* group. *Oil Gas Sci. Technol.* **2003**, *58*, 489–495. [[CrossRef](#)]
8. Müller, R.H.; Rohwerder, T.; Harms, H. Degradation of fuel oxygenates and their main intermediates by *Aquicola tertiaricarbonis* L108. *Microbiology* **2008**, *154*, 1414–1421. [[CrossRef](#)] [[PubMed](#)]
9. Auffret, M.; Labbé, D.; Thouand, G.; Greer, C.W.; Fayolle-Guichard, F. Degradation of a mixture of hydrocarbons, gasoline, and diesel oil additives by *Rhodococcus aetherivorans* and *Rhodococcus wratislaviensis*. *Appl. Environ. Microbiol.* **2009**, *75*, 7774–7782. [[CrossRef](#)]
10. McKelvie, J.R.; Hyman, M.R.; Elsner, M.; Smith, C.; Aslett, D.M.; Lacrampe-Couloume, G.; Lollar, B.S. Isotopic fractionation of methyl tert-butyl ether suggests different initial reaction mechanisms during aerobic biodegradation. *Environ. Sci. Technol.* **2009**, *43*, 2793–2799. [[CrossRef](#)]
11. Le Digabel, Y.; Demanèche, S.; Benoit, Y.; Vogel, T.M.; Fayolle-Guichard, F. Ethyl tert-butyl ether (ETBE) biodegradation by a syntrophic association of *Rhodococcus* sp. IFP 2042 and *Bradyrhizobium* sp. IFP 2049 isolated from a polluted aquifer. *Appl. Microbiol. Biot.* **2013**, *97*, 10531–10539. [[CrossRef](#)]
12. Le Digabel, Y.; Demanèche, S.; Benoit, Y.; Fayolle-Guichard, F.; Vogel, T.M. Ethyl tert-butyl ether (ETBE)-degrading microbial communities in enrichments from polluted environments. *J. Hazard. Mater.* **2014**, *279*, 502–510. [[CrossRef](#)] [[PubMed](#)]
13. Bombach, P.; Nägele, N.; Rosell, M.; Richnow, H.H.; Fischer, A. Evaluation of ethyl tert-butyl ether biodegradation in a contaminated aquifer by compound-specific isotope analysis and in situ microcosms. *J. Hazard. Mater.* **2015**, *286*, 100–106. [[CrossRef](#)]
14. Van der Waals, M.J.; Pijls, C.; Sinke, A.J.; Langenhoff, A.A.; Smidt, H.; Gerritse, J. Anaerobic degradation of a mixture of MtBE, EtBE, TBA, and benzene under different redox conditions. *Appl. Microbiol. Biot.* **2018**, *102*, 3387–3397. [[CrossRef](#)]
15. Chauvaux, S.; Chevalier, F.; Le Dantec, C.; Fayolle, F.; Miras, I.; Kunst, F.; Béguin, P. Cloning of a genetically unstable cytochrome P-450 gene cluster involved in degradation of the pollutant ethyl tert-butyl ether by *Rhodococcus ruber*. *J. Bacteriol.* **2001**, *183*, 6551–6557. [[CrossRef](#)]
16. Malandain, C.; Fayolle-Guichard, F.; Vogel, T.M. Cytochromes P450-mediated degradation of fuel oxygenates by environmental isolates. *FEMS Microbiol. Ecol.* **2010**, *72*, 289–296. [[CrossRef](#)] [[PubMed](#)]
17. Mikesková, H.; Novotný, Č.; Svobodová, K. Interspecific interactions in mixed microbial cultures in a biodegradation perspective. *Appl. Microbiol. Biot.* **2012**, *95*, 861–870. [[CrossRef](#)] [[PubMed](#)]
18. Schmidt, T.C.; Schirmer, M.; Weiss, H.; Haderlein, S.B. Microbial degradation of methyl tert-butyl ether and tert-butyl alcohol in the subsurface. *J. Contam. Hydrol.* **2004**, *70*, 173–203. [[CrossRef](#)] [[PubMed](#)]
19. Babé, A.; Labbé, D.; Monot, F.; Greer, C.; Fayolle-Guichard, F. Biodegradability of Oxygenates by Microflora from MTBE-Contaminated Sites: New Molecular Tools. In *Fuel Oxygenates. The Handbook of Environmental Chemistry*; Barceló, D., Ed.; Springer: Berlin/Heidelberg, Germany, 2007.
20. Lin, C.W.; Tsai, S.L.; Hou, S.N. Effects of environmental settings on MTBE removal for a mixed culture and its monoculture isolation. *Appl. Microbiol. Biot.* **2007**, *74*, 194–201. [[CrossRef](#)]
21. Kharoune, M.; Kharoune, L.; Lebeault, J.M.; Pauss, A. Aerobic degradation of ethyl-tert-butyl ether by a microbial consortium: Selection and evaluation of biodegradation ability. *Environ. Toxicol. Chem.* **2002**, *21*, 2052–2058. [[CrossRef](#)]
22. Van der Waals, M.J.; Plugge, C.; Meima-Franke, M.; de Waard, P.; Bodelier, P.L.; Smidt, H.; Gerritse, J. Ethyl tert-butyl ether (EtBE) degradation by an algal-bacterial culture obtained from contaminated groundwater. *Water Res.* **2019**, *148*, 314–323. [[CrossRef](#)]
23. Piveteau, P.; Fayolle, F.; Vandecasteele, J.P.; Monot, F. Biodegradation of tert-butyl alcohol and related xenobiotics by a methylotrophic bacterial isolate. *Appl. Microbiol. Biot.* **2001**, *55*, 369–373. [[CrossRef](#)] [[PubMed](#)]
24. Pospiech, A.; Neumann, B. A versatile quick-prep of genomic DNA from gram-positive bacteria. *Trends Genet.* **1995**, *66*, 217–218. [[CrossRef](#)]

25. Bruce, K.D.; Hiorns, W.D.; Hobman, J.L.; Osborn, A.M.; Strike, P.; Ritchie, D.A. Amplification of DNA from native populations of soil bacteria by using the polymerase chain-reaction. *Appl. Environ. Microbiol.* **1992**, *58*, 3413–3416. [[PubMed](#)]
26. Sanguin, H.; Herrera, A.; Oger-Desfeux, C.; Dechesne, A.; Simonet, P.; Navarro, E.; Vogel, T.M.; Moëgne-Loccoz, Y.; Nesme, X.; Grundmann, G.L. Development and validation of a prototype 16S rRNA-based taxonomic microarray for *Alphaproteobacteria*. *Environ. Microbiol.* **2006**, *8*, 289–307. [[CrossRef](#)]
27. Huber, T.; Faulkner, G.; Hugenholtz, P. Bellerophon: A program to detect chimeric sequences in multiple sequence alignments. *Bioinformatics* **2004**, *20*, 2317–2319. [[CrossRef](#)] [[PubMed](#)]
28. Wang, Q.; Garrity, G.M.; Tiedje, J.M.; Cole, J.R. Naive bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Appl. Environ. Microbiol.* **2007**, *73*, 5261–5267. [[CrossRef](#)]
29. Tamura, K.; Dudley, J.; Nei, M.; Kumar, S. MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) Software Version 4.0. *Mol. Biol. Evol.* **2007**, *24*, 1596–1599. [[CrossRef](#)]
30. Kimura, M. A simple method for estimating evolutionary rate of base substitutions through comparative studies of nucleotide sequences. *J. Mol. Evol.* **1980**, *16*, 111–120. [[CrossRef](#)]
31. Saitou, N.; Nei, M. The neighbor-joining method: A new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.* **1987**, *4*, 406–425.
32. Pommier, T.; Canback, B.; Lundberg, P.; Hagström, A.; Tunlid, A. RAMI: A tool for identification and characterization of phylogenetic clusters in microbial communities. *Bioinformatics* **2009**, *25*, 736–742. [[CrossRef](#)]
33. Kyselková, M.; Kopecký, J.; Frapolli, M.; Défago, G.; Ságová-Marečková, M.; Grundmann, G.L.; Moëgne-Loccoz, Y. Comparison of rhizobacterial community composition in soil suppressive or conducive to tobacco black root rot disease. *ISME J.* **2009**, *3*, 1127–1138. [[CrossRef](#)] [[PubMed](#)]
34. Ludwig, W.; Strunk, O.; Westram, R.; Richter, L.; Meier, H.; Yadhukumar Buchner, A.; Lai, T.; Steppi, S.; Jobb, G.; Förster, W.; et al. ARB: A software environment for sequence data. *Nucl. Acids Res.* **2004**, *32*, 1363–1371. [[CrossRef](#)]
35. Sanguin, H.; Remenant, B.; Dechesne, A.; Thioulouse, J.; Vogel, T.M.; Nesme, X.; Moëgne-Loccoz, Y.; Grundmann, G.L. Potential of a 16S rRNA-based taxonomic microarray for analyzing the rhizosphere effects of maize on *Agrobacterium* spp. and bacterial communities. *Appl. Environ. Microbiol.* **2006**, *72*, 4302–4312. [[CrossRef](#)] [[PubMed](#)]
36. Pruesse, E.; Quast, C.; Knittel, K.; Fuchs, B.M.; Ludwig, W.; Peplies, J.; Gloeckner, F.O. SILVA: A comprehensive online resource for quality checked and aligned ribosomal RNA sequence data compatible with ARB. *Nucl. Acids Res.* **2007**, *35*, 7188–7196. [[CrossRef](#)] [[PubMed](#)]
37. Stralis-Pavese, N.; Sessitsch, A.; Weilharter, A.; Reichenauer, T.; Riesing, J.; Csontos, J.; Murrell, J.C.; Bodrossy, L. Optimization of diagnostic microarray for application in analysing landfill methanotroph communities under different plant covers. *Environ. Microbiol.* **2004**, *6*, 347–363. [[CrossRef](#)] [[PubMed](#)]
38. Bouffaud, M.L.; Kyselková, M.; Gouesnard, B.; Grundmann, G.; Muller, D.; Moëgne-Loccoz, Y. Is diversification history of maize influencing selection of soil bacteria by roots? *Mol. Ecol.* **2012**, *21*, 195–206. [[CrossRef](#)] [[PubMed](#)]
39. Clarke, K.R.; Gorley, R.N. *PRIMER V6: User Manual/Tutorial, PRIMER-E Plymouth*; Plymouth Marine Laboratory: Plymouth, UK, 2006.
40. Clarke, K.R. Non-parametric multivariate analyses of changes in community structure. *Aust. J. Ecol.* **1993**, *18*, 117–143. [[CrossRef](#)]
41. Fahy, A.; McGenity, T.J.; Timmis, K.N.; Ball, A.S. Heterogeneous aerobic benzene-degrading communities in oxygen-depleted groundwaters. *FEMS Microbiol. Ecol.* **2006**, *58*, 260–270. [[CrossRef](#)] [[PubMed](#)]
42. Kyselková, M.; Kopecký, J.; Ságová-Marečková, M.; Grundmann, G.L.; Moëgne-Loccoz, Y. Oligonucleotide microarray methodology for taxonomic and functional monitoring of microbial community composition. *Plant Soil Environ.* **2009**, *55*, 379–388. [[CrossRef](#)]
43. Hendrickx, B.; Dejonghe, W.; Boëgne, M.; Brennerova, M.; Cernik, M.; Lederer, T.; Bucheli-Witschel, M.; Bastiaens, L.; Verstraete, W.; Top, E.M.; et al. Dynamics of an oligotrophic bacterial aquifer community during contact with a groundwater plume contaminated with benzene toluene ethylbenzene and xylenes: An in situ mesocosm study. *Appl. Environ. Microbiol.* **2005**, *71*, 3815–3825. [[CrossRef](#)]

44. Fahy, A.; Lethbridge, G.; Earle, R.; Ball, A.S.; Timmis, K.N.; McGenity, T.J. Effects of long-term benzene pollution on bacterial diversity and community structure in groundwater. *Environ. Microbiol.* **2005**, *7*, 1192–1199. [[CrossRef](#)] [[PubMed](#)]
45. Bundy, J.G.; Paton, G.I.; Campbell, C.D. Microbial communities in different soil types do not converge after diesel contamination. *J. Appl. Microbiol.* **2002**, *92*, 276–288. [[CrossRef](#)] [[PubMed](#)]
46. Auffret, M.D.; Yergeau, E.; Labbé, D.; Fayolle-Guichard, F.; Greer, C.W. Importance of *Rhodococcus* strains in a bacterial consortium degrading a mixture of hydrocarbons, gasoline, and diesel oil additives revealed by metatranscriptomic analysis. *Appl. Microbiol. Biotechnol.* **2015**, *99*, 2419–2430. [[CrossRef](#)] [[PubMed](#)]
47. Hatzinger, P.B.; McClay, K.; Vainberg, S.; Tugusheva, M.; Condee, C.W.; Steffan, R.J. Biodegradation of methyl tert-butyl ether by a pure bacterial culture. *Appl. Environ. Microbiol.* **2001**, *67*, 5601–5607. [[CrossRef](#)] [[PubMed](#)]
48. Raynal, M.; Pruden, A. Aerobic MTBE biodegradation in the presence of BTEX by two consortia under batch and semi-batch conditions. *Biodegradation* **2008**, *19*, 269–282. [[CrossRef](#)]
49. Bastida, F.; Rosell, M.; Franchini, A.G.; Seifert, J.; Finsterbusch, S.; Jehmlich, N.; Jechalke, S.; von Bergen, M.; Richnow, H.H. Elucidating MTBE degradation in a mixed consortium using a multidisciplinary approach. *FEMS Microbiol. Ecol.* **2010**, *73*, 370–384. [[CrossRef](#)] [[PubMed](#)]
50. Schuster, J.; Purswani, J.; Breuer, U.; Pozo, C.; Harms, H.; Mi, R.H.; Rohwerder, T. Constitutive expression of the cytochrome P450 EthABCD monooxygenase system enables degradation of synthetic dialkyl ethers in *Aquicola tertiarycarbonis* L108. *Appl. Environ. Microbiol.* **2013**, *79*, 2321–2327. [[CrossRef](#)]



© 2019 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).